



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

great handfulls of the flowers. I think they must have opened about a week ago, as the racemes had set fruit below. There were many budding clusters also. I account for this display by the fact that the swamp in question was reduced to dusty dryness by the long-continued drouth; this acted on the plants much as their normal winter rest. When followed by rain—flooding the marsh—and then by an extended period of warm weather, they burst forth into flower. A natural consequence, I should suppose, would be a dearth of my favorites next spring.—W. WHITMAN BAILEY, *Brown University*.

Tubers.—That the scales on the tubers represent leaves of the of aerial stems is well known. The study of the phyllotaxy of these subterranean leaves is quite as interesting as that of ordinary leaves. In examining all tubers of cultivated and wild plants that I can obtain, I find that a plant has the same arrangement of foliar organs on tubers that it has on the stem; and where two plans exist, the one at the base of the stem is the one followed by the tuber. Good examples are found in the potato, in which both leaves and scales are alternate; and in *Helianthus doricoides*, L., and *tuberosus*, L., where the leaves are opposite below and more or less alternate above, the scales on the tubers are opposite. In the latter species scales frequently subtend “knobs”, the tuber branches, which are then opposite and themselves bear scales—the leaves of the branch. The dimerous whorls decussate on tubers as well as do those of the stem. Another interesting fact is the completion of growth as to length in the lower internodes of the tuber while the upper are still quite small—a characteristic of the stem.—AUG. F. FOERSTE, *Dayton, Ohio*.

Notes.—Teachers of botany may be interested in knowing, if they do not know already, that the now common Japanese *Ampelopsis* presents an excellent instance of a uni-foliate compound leaf. The three-lobed, or sometimes barely lobed and dentate leaflet has all the appearance of a simple leaf, but falls off by a distinct articulation from the top of the extremely long petiole, which is apt to persist sometime thereafter.

In the analysis of *Heterocentron roseum*, of the order *Melastomaceæ*, students complain that they can not ascertain the name by the key in the School and Field-book of Botany. I find that the difficulty is in the statement that in that family the calyx is coherent with the ovary. In *Heterocentron*, so far as I have examined specimens, it is distinctly *free*. The ordinal characteristics given by LeMaout and Decaisne, give the alternative of free or coherent.—W. W. BAILEY, *Brown University*.